



## Evaluation of Efficacy of Perioperative Oral Dexamethasone in Reducing Swelling and Ecchymosis After Blepharoplasty Surgery

Rabar Abas Hama amin\* Zanyar Mustafa Amin\*\* Bayad Jaza Mahmood\*\*\*

---

### Abstract

**Background and objectives:** Blepharoplasty, a common cosmetic procedure, can lead to postoperative issues like swelling and bruising. Steroids are occasionally used to address these concerns and speed up recovery. This study aims to evaluate how effective steroids are in reducing periorbital swelling and bruising in blepharoplasty patients.

**Methods:** From November 2022 to June 2023, a descriptive analysis was conducted on forty participants at Dermodonto Center and Safin private hospitals in Kurdistan/Iraq. The participants were divided into two groups. Group 1 (twenty individuals) received Dexamethasone 8 mg tablets thirty minutes before surgery, continuing for three days post-surgery. Group 2 (twenty participants) did not receive steroid treatment. Periorbital swelling and bruising were assessed using a standardized visual inspection scale, enabling objective measurement and comparison between the steroid-treated and non-treated groups.

**Results:** Among the 40 patients, 35 were female and 5 were male, with an average age of 39.19. Median swelling scores decreased from 1.5 (range: 1.0-2.0) on POD 1 to 0 (range: 0.0-1.0) on POD 7. Similarly, median ecchymosis scores decreased from 2.0 (range: 1.0-2.0) on POD 1 to 0 (range: 0.0-1.0) on POD 7. The study demonstrated a significant reduction in swelling and ecchymosis among participants who received steroid treatment compared to the control group, with statistically significant results ( $p < 0.05$ ).

**Conclusion:** Steroids reduce periorbital swelling and bruising after blepharoplasty, enhancing outcomes and patient satisfaction, and highlighting their procedural benefits.

**Keywords:** Blepharoplasty, Dexamethasone, Ecchymosis, Edema

---

\*B.D.S Candidate of Kurdistan Higher Council of Medical Specialties / Oral and Maxillofacial Surgery/

Corresponding author: Email: [rabar.abass@yahoo.com](mailto:rabar.abass@yahoo.com)

\*\*B.D.S. FIBMS. / Assistant Professor at Kurdistan Higher Council of Medical Specialties, Director of the Maxillofacial Center/ Sulaimani/ Email: [zanyar.amin@univsul.edu.iq](mailto:zanyar.amin@univsul.edu.iq)

\*\*\*B.D.S. FKBMS. AACME. Assistant Professor at Ministry of Higher Education, Head of Oral and Maxillofacial Surgery Department/ University of Sulaimani/ College of Dentistry. Email: [bayad.mahmood@univsul.edu.iq](mailto:bayad.mahmood@univsul.edu.iq)



## Introduction

In the world of cosmetic surgery, procedures such as blepharoplasty are undergoing a substantial surge in popularity. This upswing in demand arises from its impressive ability not only to rectify visible cosmetic flaws but also to notably enhance the overall facial appearance. Beyond simply improving looks, blepharoplasty can also enhance facial symmetry and give a more youthful and vibrant look. Its ability to deliver such comprehensive results has caught the attention of many people looking for a complete facial rejuvenation, making blepharoplasty a highly sought-after choice in the field of cosmetic enhancement.<sup>1</sup> Blepharoplasty, despite its potential to enhance facial aesthetics, presents its own set of challenges. Following surgery, patients frequently encounter discomfort, swelling, and bruising. The severity and duration of these complications hinge on various factors, including the complexity of the surgical procedure and each individual's unique healing responses. Consequently, effectively managing these postoperative issues is imperative to ensure optimal patient outcomes and satisfaction, underscoring the importance of comprehensive care throughout the recovery process.<sup>2-4</sup> Furthermore, when multiple facial procedures are performed simultaneously, like combining rhinoplasty with blepharoplasty, the chances and severity of postoperative swelling and bruising increase. This highlights the importance of thorough preoperative assessment and patient education. It's crucial to manage expectations and minimize possible complications associated with combined procedures to achieve the best results and ensure patient satisfaction. Therefore, conducting a comprehensive evaluation and maintaining transparent communication with patients are vital steps to prepare them for potential challenges and ensure they are fully informed

and engaged in their treatment process.<sup>5</sup> Over the course of medical history, corticosteroids have played a crucial part in managing postoperative bruising and swelling, especially in complex maxillofacial surgeries. Leveraging their powerful anti-inflammatory properties, steroids have shown significant effectiveness in reducing tissue swelling and inflammation. This therapeutic ability has been invaluable in speeding up recovery and easing discomfort for patients undergoing intricate facial procedures. Consequently, the strategic use of corticosteroids in perioperative care routines has become fundamental in improving patient outcomes and enriching the overall surgical experience for complex facial surgeries.<sup>3-9</sup> In the field of Cranio-Facio-Maxillary surgeries, steroids are commonly used perioperatively to reduce swelling after surgery and accelerate the healing process. Several reports and studies have emphasized the effectiveness of systemic steroid administration, as well as other agents like Bromelain and topical products such as arnica, in minimizing facial swelling following various procedures. These procedures include orbital floor exploration, bimaxillary osteotomy, third molar extraction, hair transplant, and rhinoplasty, among others. Such research highlights the versatility and efficacy of steroids in managing postoperative swelling across a range of craniofacial surgeries, providing valuable insights for improving patient outcomes in these intricate surgical settings.<sup>4,5,10</sup> In consideration of this background, our research was designed to meticulously investigate the role of corticosteroids in addressing the occurrence of postoperative bruising and swelling after blepharoplasty surgery. Our study's focal point on this specific aspect of perioperative care was intended to offer a comprehensive understanding and provide actionable insights aimed at refining patient outcomes





and enhancing the overall journey for individuals undergoing blepharoplasty procedures. Through a thorough exploration of the efficacy of corticosteroids in this context, we aimed to contribute substantively to the advancement of clinical practices in the field of cosmetic surgery, ultimately striving to elevate the standard of care and satisfaction for patients undergoing blepharoplasty.

### Patients and methods:

The research involved 40 randomly selected healthy adult individuals who sought blepharoplasty surgery treated at Dermodonto Center and Safin private hospitals in Kurdistan/Iraq. The patients represented wide range of demographics, including an age, gender distribution, diverse socioeconomic backgrounds, occupations and comprehensive recording of the patient's data after operations. After receiving comprehensive information about the procedure, as well as the benefits and potential risks associated with steroid use, the patients are provided explicit consent for both the surgical intervention and the administration of steroids. Exclusion criteria included systemic disorders, history of chemotherapy or radiation therapy, pregnancy or lactation, smoking, wound healing therapies, anticoagulant drugs, and hypersensitivity. Patients underwent blepharoplasty surgery performed by the same author under local anesthesia with 1-3 dental carpules (1.8ml) containing a 1:100,000 epinephrine solution. Patients were divided into two groups: the treatment group (n = 20) received dexamethasone 8mg tablet one day before and half an hour before the procedure, continuing for 3 days postoperatively, while the control group (n = 20) did not receive perioperative steroids. Patient scores for ecchymosis and swelling were recorded and assessed at the same surgical clinic using a visual inspection scale by three different observers on postoperative

days 1, 3, and 7, following the Totonchi method.<sup>11</sup> For ecchymosis, grading ranged from 0 to 4: 0 represented no color change, 1 indicated a yellowish color, 2 denoted a light purple color, 3 signified a dark purple color, and 4 represented a very dark purple color. Meanwhile, the swelling was graded on a scale of 0 to 4: 0 indicated no edema, 1 denoted minimal edema, 2 indicated swelling covering the iris, 3 represented swelling extending to the pupil, and 4 indicated massive edema. With the accordance of Helsinki Declaration, the study was approved by the ethical committee of Kurdistan Higher Council of Medical Specialties (Reference number 844, dated April 12, 2023). Data was entered using IBM SPSS version 22 software. Ecchymosis and swelling were recorded and assessed using a visual inspection scale on postoperative days 1, 3, and 7. Gender differences were analyzed using Pearson's chi-square test, and continuous variables between groups were compared using Kruskal Wallis analysis of variance.

### Results

Of those 40 patients who underwent a blepharoplasty operation, 35 were women and the other 5 were men. The patients were between the ages of 25 and 65, with an average age of 39.19. The study included a treatment group (20 patients) receiving steroid injections and a control group (20 patients) without receiving the injections. Regarding age, gender, and the duration of operation, there was not a statistically significant difference between the treatment and control groups. Neither the operation nor the use of steroids caused any negative effects. The mean score of swelling and ecchymosis at postoperative days 1 and 3 was significantly lower in the group that received dexamethasone treatment than that of the control group. The edema was still diminished at postoperative day 7. However, this fast reduction in the swelling and ecchymosis was not seen in the control



group. Postoperative edema and ecchymosis were significantly reduced in the group receiving dexamethasone on postoperative days 1, 3, and 7 with a statistically significant difference (P value <0.001). The surgical outcomes regarding edema and ecchymosis are summarized in Table (1) and illustrated in Figures (1, and 2).


**Table (1):** Median value of swelling and ecchymosis in study groups:

| Variable   | Treatment group<br>(n=20) | Control group<br>(n=20) | p value |
|------------|---------------------------|-------------------------|---------|
| Swelling   |                           |                         |         |
| POD* 1     | 1.5 (1.0-2.0)             | 3 (2.75-4.0)            | < 0.001 |
| POD 3      | 1 (1.0-1.0)               | 2.5 (1.0-3.0)           | < 0.001 |
| POD 7      | 0 (0.0-1.0)               | 1 (1.0-2.0)             | < 0.004 |
| Ecchymosis |                           |                         |         |
| POD 1      | 2 (1.0-2.0)               | 3.5 (3.0-4.0)           | < 0.001 |
| POD 3      | 1 (1.0-2.0)               | 3 (2.0-3.25)            | < 0.001 |
| POD 7      | 0 (0.0-1.0)               | 1 (1.0-2.0)             | 0.013   |

POD\* =Post Operative Day



**Figure (1):** A patient in the control group, from above to below, A=1st, B=3rd, and C=7th postoperative day.



**Figure (2):** A patient in the treatment group, from above to below, A=1st, B=3rd, and C=7th postoperative day.





## Discussion

The Glucocorticoids' anti-inflammatory properties reduce vascular permeability, resulting in reduced exudates and edema. Dexamethasone is a very effective anti-inflammatory steroid having a biological half-life of 36-45 hours.<sup>2-4</sup> Recently, many surgeons have advocated for reducing postoperative bruising and swelling following cosmetic surgery. To address this concern, steroids are administered in different doses both before and after the surgical procedure.<sup>4-9</sup> In this study, a specific grading system was devised to evaluate ecchymosis severity and edema degree. Consistent with prior research, our findings revealed reduced peri-orbital edema scores on postoperative days 1 and 4 among patients administered dexamethasone, underscoring its anti-edema efficacy in the early postoperative phase. Nevertheless, it is crucial to note that potential adverse effects were associated with long-term steroid use.<sup>7,12</sup> Long-term steroid usage can result in mineral deficiencies, edema, coronary artery disease, hypertension, Diabetes, and renal failure in addition to infection. Gastric ulcers and pancreatitis are gastrointestinal complications; skin atrophy, pimples, hirsuteness hair loss, and tendency to get bruising are dermatologic side consequences; osteoporosis, osteonecrosis, and myopathy are musculoskeletal complications; and cataracts and myopia are ophthalmological complications.<sup>12-16</sup> However, for those who do not have particular contraindications, short-term corticosteroid treatment usually has few side effects. Edema and ecchymosis commonly occur as side effects of facial surgery. To counteract these effects, varying doses of steroids were administered before, during, and after the surgical procedure.<sup>16,17</sup> The majority of studies revealed that steroids minimize the amount of swelling and ecchymosis in the facial region.<sup>6-11,18-22</sup> A study observed that steroids reduced facial

edema in 158 individuals who underwent various facial procedures after administering a single bolus (1g dosage) of methylprednisolone. In the field of faciomaxillary surgery<sup>17</sup>, the use of steroids is approved for the surgical extraction of third molars and facial osteotomies.<sup>2,3,22</sup> In a study of third molar surgery, one dose of 125 mg methylprednisolone was found to reduce trismus, edema, and face pain.<sup>18</sup> Similarly, another author investigated the effects of one dose each of 1.5 mg/kg and 3 mg/kg of methylprednisolone following mandibular third molar surgery. Their study revealed that steroids effectively reduced both pain and facial swelling, with no significant advantage observed for the higher dose (3 mg/kg) compared to the lower dose (1.5 mg/kg).<sup>21</sup> In a study involving 40 patients undergoing rhinoplasty with osteotomy, Gurlek, administered high doses of methylprednisolone (250 and 500 mg). Their findings revealed that a larger dose of methylprednisolone was effective in avoiding and lowering periorbital edema after open rhinoplasty with osteotomy.<sup>5</sup> In another study involving 42 patients undergoing open rhinoplasty, the statistical analysis of patient outcomes indicates that a suprapariosteal injection of dexamethasone did not effectively reduce swelling and bruising after rhinoplasty.<sup>9</sup> Daniel et al. demonstrated that short-term steroid administration effectively reduces periorbital swelling after orbital surgery.<sup>17</sup> Palpebral edema after rhinoplasty may impede vision during the initial postoperative period. Furthermore, postoperative peri-orbital edema and ecchymosis might cause discomfort and anxiety.<sup>10</sup> In a randomized trial with 84 open rhinoplasty patients, two doses of 10 mg dexamethasone were given during and after surgery. Results showed reduced periorbital swelling, bruising, and intraoperative bleeding compared to a placebo, possibly due to the





interaction between steroids and adrenaline.<sup>23</sup> Bian, X. et al, in a systematic review of ten articles involving 374 participants support using dexamethasone during rhinoplasty to reduce intraoperative blood loss, edema, and ecchymosis compared to normal saline. Meta-analysis indicates significant benefits, with medium to large effect reductions in these parameters compared to placebo groups receiving normal saline.<sup>24</sup> The present study demonstrated statistically significant differences in postoperative swelling and ecchymosis between the treatment and control groups, highlighting the efficacy of dexamethasone.

### Limitations:

However: the sample size is relatively small, which may limit the generalizability of the findings. Larger studies are needed to confirm these results. And The study population consisted predominantly of women, which may limit the applicability of the findings to male patients or more diverse demographic groups.

### Conclusion

The study demonstrates differences between the group who received steroids and the control group that are statistically and clinically significant. Periorbital ecchymosis and edema can be prevented and reduced with the use of preoperative steroid injection.

### Recommendations

Future studies should include a larger number of participants to enhance the statistical power and generalizability of the findings.

### Acknowledgment:

The authors would like to acknowledge the Oral and Maxillofacial Department, University of Sulaimani, and Kurdistan Higher Council of Medical Specialties for supporting the study.

### Conflict of Interest:

There is no conflict of interest to declare.

### References:

1. Defraia B., Focardi M., Grassi S., Chiavacci G., Faccioli S., Romano G., et al. Negative Outcomes of Blepharoplasty and Thyroid Disorders: Is Compensation Always Due? A Case Report with a Literature Review. *Diseases*. 2024 Apr 10;12(4):75.
2. Fatah G., Mahmood B. Preoperative submucosal dexamethasone injection for minimizing postoperative pain, trismus, and edema associated with impacted mandibular third molar surgery. *J Craniomaxillofac Res* 2022; 9(3): 132-138.
3. Bakri M., Alabdali F., Mahzari R., Rajhi T., Gohal N., Sufyani R., et al. Comparison of the effects of two different styles of orally prescribing prednisolone on postoperative sequelae of surgical extraction of an impacted mandibular third molar: a single-blind randomized study. *J Korean Assoc Oral Maxillofac Surg*. 2024 Feb 29;50(1):27-34.
4. Mahmood B. The effect of using pineapple fresh juice to improve post-surgical pain, ecchymosis, and swelling in the maxillofacial region. *J Craniomaxillofac Res*. 2021;8(2):67-75.
5. Gürlek A., Fariz A., Aydoğan H., Ersöz A., Evans G. Effects of high dose corticosteroids in open rhinoplasty. *J Plast Reconstr Aesthet Surg*. 2009 May;62(5):650-5.
6. Valente D., Steffen N., Carvalho L., Borille G., Zanella R., Padoin, A. Preoperative Use of Dexamethasone in Rhinoplasty: A Randomized, Double-blind, Placebo-Controlled Clinical Trial. *JAMA Facial Plast Surg*. 2015 May-Jun;17(3):169-73.
7. Tuncel U., Turan A., Bayraktar M., Erkorkmaz U., Kostakoglu N. Efficacy of dexamethasone with controlled hypotension on intraoperative bleeding,





- postoperative oedema and ecchymosis in rhinoplasty. *J Craniomaxillofac Surg*. 2013 Mar;41(2):124-8.
8. Coroneos C., Voineskos S., Cook D., Farrokyar F., Thoma, A. Perioperative Corticosteroids Reduce Short-Term Edema and Ecchymosis in Rhinoplasty: A Meta-Analysis. *Aesthet Surg J*. 2016 Feb;36(2):136-46.
  9. Jafari M., Maleki D., Jahandideh H., Rajaeih S., Ghashghaei S., Wood D. The Effect of Subcutaneous Dexamethasone to Reduce Edema and Ecchymosis in Rhinoplasty Patients. *Int J Otolaryngol*. 2022 Feb 14; 2022:3054767.
  10. Simsek G., Sari E., Kilic R., Bayar M. Topical Application of Arnica and Mucopolysaccharide Polysulfate Attenuates Periorbital Edema and Ecchymosis in Open Rhinoplasty: A Randomized Controlled Clinical Study. *Plast Reconstr Surg*. 2016 Mar;137(3): e530-535.
  11. Totonchi A., Guyuron B. A randomized, controlled comparison between arnica and steroids in the management of post rhinoplasty ecchymosis and edema. *Plast Reconstr Surg*. 2007 Jul;120(1):271-274.
  12. Malhari A., Sharma R., Rattan V. Role of corticosteroids in reduction of postoperative edema in craniofaciomaxillary surgery. *J. Evolution Med. Dent. Sci*. 2016;5(48):3090-3103.
  13. Tamene S., Dalhoff K., Schwarz P., Backer V., Aanaes, K. Systemic corticosteroids in treatment of chronic rhinosinusitis-A systematic review. *Eur Clin Respir J*. 2023 Aug 4;10(1):2240511.
  14. Duru N., Van Der Goes M., Jacobs J., Andrews T., Boers M., Buttgereit F., et al. EULAR evidence-based and consensus-based recommendations on the management of medium to high-dose glucocorticoid therapy in rheumatic diseases. *Ann Rheum Dis*. 2013 Dec;72(12):1905-13.
  15. He J., He S., Zhang Y., Tian Y., Hao P., Li T., et al. Association between intraoperative steroid and postoperative mortality in patients undergoing craniotomy for brain tumor. *Front Neurol*. 2023 Jun 29; 14:1153392.
  16. Abdu S., Alsubhi A., Alzahrani A., Alshehri A., Altaylouni N., Rammal, A. Comparison of oral versus intravenous steroid in the management of Bell's palsy: a systematic review and meta-analysis of randomized clinical trials. *Eur Arch Otorhinolaryngol*. 2024 Mar;281(3):1095-1104.
  17. Daniel A., Warren A., Jessica N., Jamal M., Larry H., Samuel S. Perioperative Steroids for Minimizing Edema and Ecchymosis After Rhinoplasty: A Meta-Analysis. *Aesth Surg J*. 2011;31(6) 648–657.
  18. Esen E., Taşar F., Akhan O. Determination of the anti-inflammatory effects of methylprednisolone on the sequelae of third molar surgery. *J Oral Maxillofac Surg*. 1999 Oct;57(10):1201-6.
  19. Geng W., Nie Y., Huang S. Effects of methylprednisolone on the duration of rocuronium-induced neuromuscular block: A randomized double-blind trial. *Med (Baltimore)*. 2017 Sep;96(39): e7947.
  20. Levin M., Ziai H., Roskies M. Modalities of Post-Rhinoplasty Edema and Ecchymosis Measurement: A Systematic Review. *Plast Surg (Oakv)*. 2022 May;30(2):164-174.
  21. Garg D. The Comparison of the effect of two different doses of methylprednisolone on pain, swelling and trismus after 3rd molar surgery. *J. Dent. Sci*. 2022;8(2):06-11.
  22. Koçer G., Yuce E., Tuzuner A., Dereci O., Koskan, O. Effect of the route of





- administration of methylprednisolone on oedema and trismus in impacted lower third molar surgery. *Int J Oral Maxillofac Surg.* 2014 May;43(5):639-43.
23. Alajmi M., Al-Abdulahdi K., Al-Noumas H., Kavitha G. Results of intravenous steroid injection on reduction of postoperative edema in rhinoplasty. *Indian J Otolaryngol Head Neck Surg.* 2009 Dec;61(4):266-9.
24. Bian X., Liu H., Sun J., Zhang X., Li N., Chen, M. Efficacy of Dexamethasone for Reducing Edema and Ecchymosis After Rhinoplasty: A Systematic Review and Meta-analysis. *Aesth Plast Surg* 2020; 44, 1672–1684.

